TABLE OF CONTENTS

HE PET	TITIONERS
NTRODU	JCTION
CTION	REQUESTED
TATEME	ENT OF GROUNDS
1. B	Senzodiazepines Should Only Be Used For Two To Four Weeks
	A. Ativan's drug literature compared to other FDA-approved
	B. Drug trials only last a few months
	C. It is well established benzodiazepines should only be
	D. International guidelines
	E. Group survey
	Risks Of Developing Physical Dependency And Withdrawal Symptoms With Short Term Benzodiazepine Use
	A. Physical dependency
	B. Tolerance/tolerance withdrawal
	C. Withdrawal
	D. Misinformation: tolerance, physical dependency, and withdrawal
3.	Protracted Withdrawal
	A. Benzodiazepines are known to have a post withdrawal syndrome
	B. Group survey
4.	Confusion Regarding The Terms "Physical Dependency" And "Addiction"
5.	Benzodiazepine Equivalencies
6.	Withdrawal Methods
	A. Gradual dose reduction is not defined
	B. Patients are often told by their doctors
	C. Group survey
	D. The following withdrawal methods, though rarely known in the medical community, are the most successful strategies
	E. The elements of a well designed taper
7.	Clear Information Needs To Get To Doctors, Pharmacists, And Patients
8.	Apply A Black Box Warning To All FDA-Approved Benzodiazepines
0.	Apply A Didek Dex training to the Data person Derizonal
ERSON	AL IMPACT STATEMENTS
PPOSIT	TION
ONCLU	ISION
	IMENTAL IMPACT STATEMENT
ERTIFIC	
OCTOR	RS AND RESEARCHERS WHO ENDORSE THIS PETITION